



For more information about **Measles** contact:

*The Community Infection Control Nurse at your
local Primary Care Trust (PCT)*

or your local Health Protection Unit

or

<http://www.hpa.org.uk/topics/index.htm>

HPA North West



<http://www.hpa.org.uk/northwest/factsheets/MEASLES.pdf>

Information produced by: Health Protection Agency North West

Date: September 2007

Review Date: September 2009

Measles

INFORMATION LEAFLET

What is measles?

Measles is an infection caused by a virus (germ) and symptoms include:

- Fever
- Irritability
- Cold-like symptoms - running nose
- Sore and runny eyes
- Dry “croupy” cough
- White spots on the gums (2nd and 3rd day)
- Rash (3rd or 4th day) starting on the face and behind the ears, then spreading down the body.
- Diarrhoea - often in the early stages in children.

Serious complications of measles infection can occur such as ear infections (approx. 1 in 20), bronchitis/pneumonia (approx. 1 in 25), convulsions (approx. 1 in 200), and in rare cases, encephalitis (infection of the brain) or meningitis (approx. 1 in 1,000) may occur. The likelihood of problems varies with age, and complications are generally more common in young babies, older children and adults.

Who does it affect?

ANYONE - who has not developed natural immunity or has not been vaccinated against measles. Children are most vulnerable though, but babies up to approximately 6 months are protected by their mother’s antibodies.

How do you get measles?

MEASLES IS VERY INFECTIOUS and you catch it by being in close contact with someone who already has the infection. The virus is passed in the secretions of the infected person’s nose and throat and is spread by their sneezing and coughing. Also it is possible to catch measles from direct contact with articles which have been contaminated by these infected secretions e.g. hankies.

How do you prevent measles?

- **By immunisation** - there is a safe and effective vaccine, which protects against measles. It is one of the “M” components in MMR

vaccine and a child needs two doses of this vaccine, one at 13 - 15 months and another dose pre-school.

- Protection can be life long.
- If a child or young adult has not received 2 doses of MMR vaccine, this can be given at any age.
- By avoiding close personal contact with a person with measles if you are at risk. Because measles vaccine is a “live” vaccine, it is not recommended that children who have a weakened immune system e.g. because of medication or chemotherapy, should be vaccinated so these children are particularly vulnerable and should avoid contact whenever possible.
- If you are pregnant and you have been in contact with measles always discuss with your GP or midwife immediately so that he/she can advise you.

Do you need to stay off school/work?

YES - MEASLES IS VERY INFECTIOUS.

Measles is infectious from just before the symptoms first occur and for approximately 4 days after the appearance of the rash, so general advice is to stay off school or work for **at least 7 days after symptoms first develop.**

The incubation period (the period when someone has the infection but is not displaying symptoms) is on average 10 days.

Do you need treatment?

There is no specific treatment for measles but an appropriate medicine to help bring down a high temperature should be used (**DO NOT USE ASPIRIN IN CHILDREN UNDER 16 YEARS OF AGE**) and if necessary, sponging down with tepid water. Also, if a secondary infection develops then the doctor may prescribe antibiotics to treat this e.g. an ear or chest infection.

2 – 6 weeks after the illness, a swab may be taken from the mouth of the infected person. This is to confirm if it was a definite case of measles. Your GP will be informed of the results.